

relations (10.235–10.237), then so do

$$\mathbf{J} \quad \text{and} \quad -\mathbf{K}. \quad (10.238)$$

The infinitesimal Lorentz transformation (10.234) is the  $4 \times 4$  matrix

$$L = I + \omega = I + \theta_\ell R_\ell + \lambda_j B_j = \begin{pmatrix} 1 & \lambda_1 & \lambda_2 & \lambda_3 \\ \lambda_1 & 1 & -\theta_3 & \theta_2 \\ \lambda_2 & \theta_3 & 1 & -\theta_1 \\ \lambda_3 & -\theta_2 & \theta_1 & 1 \end{pmatrix}. \quad (10.239)$$

It moves any 4-vector  $x$  to  $x' = Lx$  or in components  $x'^a = L^a_b x^b$

$$\begin{aligned} x'^0 &= x^0 + \lambda_1 x^1 + \lambda_2 x^2 + \lambda_3 x^3 \\ x'^1 &= \lambda_1 x^0 + x^1 - \theta_3 x^2 + \theta_2 x^3 \\ x'^2 &= \lambda_2 x^0 + \theta_3 x^1 + x^2 - \theta_1 x^3 \\ x'^3 &= \lambda_3 x^0 - \theta_2 x^1 + \theta_1 x^2 + x^3. \end{aligned} \quad (10.240)$$

More succinctly with  $t = x^0$ , this is

$$\begin{aligned} t' &= t + \boldsymbol{\lambda} \cdot \mathbf{x} \\ \mathbf{x}' &= \mathbf{x} + t\boldsymbol{\lambda} + \boldsymbol{\theta} \wedge \mathbf{x} \end{aligned} \quad (10.241)$$

in which  $\wedge \equiv \times$  means cross-product.

For arbitrary real  $\boldsymbol{\theta}$  and  $\boldsymbol{\lambda}$ , the matrices

$$L = e^{-i\theta_\ell J_\ell - i\lambda_\ell K_\ell} \quad (10.242)$$

form the subgroup of  $SO(3,1)$  that is connected to the identity matrix  $I$ . This subgroup preserves the sign of the time of any time-like vector, that is, if  $x^2 < 0$ , and  $y = Lx$ , then  $y^0 x^0 > 0$ . It is called the proper orthochronous Lorentz group. The rest of the (homogeneous) Lorentz group can be obtained from it by space  $\mathcal{P}$ , time  $\mathcal{T}$ , and space-time  $\mathcal{PT}$  reflections.

The task of finding all the finite-dimensional irreducible representations of the proper orthochronous homogeneous Lorentz group becomes vastly simpler when we write the commutation relations (10.235–10.237) in terms of the hermitian matrices

$$J_\ell^\pm = \frac{1}{2} (J_\ell \pm iK_\ell) \quad (10.243)$$

which generate two independent rotation groups

$$\begin{aligned} [J_i^+, J_j^+] &= i\epsilon_{ijk} J_k^+ \\ [J_i^-, J_j^-] &= i\epsilon_{ijk} J_k^- \\ [J_i^+, J_j^-] &= 0. \end{aligned} \quad (10.244)$$

Thus the Lie algebra of the Lorentz group is equivalent to two copies of the Lie algebra (10.100) of  $SU(2)$ . Its finite-dimensional irreducible representations are the direct products

$$D^{(j,j')}(\boldsymbol{\theta}, \boldsymbol{\lambda}) = e^{-i\theta_\ell J_\ell - i\lambda_\ell K_\ell} = e^{(-i\theta_\ell - \lambda_\ell)J_\ell^+} e^{(-i\theta_\ell + \lambda_\ell)J_\ell^-} \quad (10.245)$$

of the nonunitary representations  $D^{(j,0)}(\boldsymbol{\theta}, \boldsymbol{\lambda}) = e^{(-i\theta_\ell - \lambda_\ell)J_\ell^+}$  and  $D^{(0,j')}(\boldsymbol{\theta}, \boldsymbol{\lambda}) = e^{(-i\theta_\ell + \lambda_\ell)J_\ell^-}$  generated by the three  $(2j+1) \times (2j+1)$  matrices  $J_\ell^+$  and by the three  $(2j'+1) \times (2j'+1)$  matrices  $J_\ell^-$ . Under a Lorentz transformation  $L$ , a field  $\psi_{m,m'}^{(j,j')}(x)$  that transforms under the  $D^{(j,j')}$  representation of the Lorentz group responds as

$$U(L) \psi_{m,m'}^{(j,j')}(x) U^{-1}(L) = D_{mm''}^{(j,0)}(L^{-1}) D_{m'm'''}^{(0,j')}(L^{-1}) \psi_{m'',m'''}^{(j,j')}(Lx). \quad (10.246)$$

Although these representations are not unitary, the  $SO(3)$  subgroup of the Lorentz group is represented unitarily by the hermitian matrices

$$\mathbf{J} = \mathbf{J}^+ + \mathbf{J}^-. \quad (10.247)$$

Thus, the representation  $D^{(j,j')}$  describes objects of the spins  $s$  that can arise from the direct product of spin- $j$  with spin- $j'$  (Weinberg, 1995, p. 231)

$$s = j + j', j + j' - 1, \dots, |j - j'|. \quad (10.248)$$

For instance,  $D^{(0,0)}$  describes a spinless field or particle, while  $D^{(1/2,0)}$  and  $D^{(0,1/2)}$  respectively describe **left**-handed and **right**-handed spin-1/2 fields or particles. The representation  $D^{(1/2,1/2)}$  describes objects of spin 1 and spin 0—the spatial and time components of a 4-vector.

The generators  $K_j$  of the Lorentz boosts are related to  $\mathbf{J}^\pm$  by

$$\mathbf{K} = -i\mathbf{J}^+ + i\mathbf{J}^- \quad (10.249)$$

which like (10.247) follows from the definition (10.243).

The interchange of  $\mathbf{J}^+$  and  $\mathbf{J}^-$  replaces the generators  $\mathbf{J}$  and  $\mathbf{K}$  with  $\mathbf{J}$  and  $-\mathbf{K}$ , a substitution that we know (10.238) is legitimate.

### 10.32 Two-Dimensional Representations of the Lorentz Group

The generators of the representation  $D^{(1/2,0)}$  with  $j = 1/2$  and  $j' = 0$  are given by (10.247 & 10.249) with  $\mathbf{J}^+ = \boldsymbol{\sigma}/2$  and  $\mathbf{J}^- = 0$ . They are

$$\mathbf{J} = \frac{1}{2}\boldsymbol{\sigma} \quad \text{and} \quad \mathbf{K} = -i\frac{1}{2}\boldsymbol{\sigma}. \quad (10.250)$$